

USDA Foreign Agricultural Service

GAIN Report

Global Agricultural Information Network

THIS REPORT CONTAINS ASSESSMENTS OF COMMODITY AND TRADE ISSUES MADE BY
USDA STAFF AND NOT NECESSARILY STATEMENTS OF OFFICIAL U.S. GOVERNMENT
POLICY

Voluntary _ Public

Date: 7/15/2011

GAIN Report Number: JA1036

Japan

Post: Tokyo

Information from Fishery Agency Japan

Report Categories:

Fishery Products

Approved By:

Steve Shnitzler

Prepared By:

Yuichi Hayashi

Report Highlights:

This is an agenda provided by Fisheries Agency Japan for interagency working group on ocean plume modeling, June 23, 2011. The title of the report is "To Secure the Safety of Fishery Products in Japan - In response discharges of radioactive substances to the sea from the TEPCO Fukushima Daiichi Nuclear Power Plant."

General Information:

June, 2011
Fisheries Agency of Japan

To Secure the Safety of Fishery Products in Japan

[In response to discharges of radioactive substances to the sea
from the TEPCO Fukushima Daiichi Nuclear Power Plant]

Summary

Radioactive substances have been detected in the seawaters in the vicinity of the TEPCO Fukushima Daiichi Nuclear Power Plant, which originated from discharges of contaminated water, atmospheric fallout and washout with precipitation into the sea.

In response to the critical situation, the Japanese government swiftly introduced monitoring programs to measure the levels of radioactive substances contained in fishery products as well as in seawaters. So far, only a small number of species caught in the area close to the Fukushima NPP show the levels of radioactive substances exceeding the Provisional Regulatory Values*. The Japanese government ensures the safety of fishery products on the market, by imposing suspension of related fishing activities and market distribution as soon as a sampling measurement of a fishery product detects a level exceeding the Provisional Regulatory Values.

[(*) Indices for restrictions on intake of foods set by the Ministry of Health, Labor and Welfare. Japan's index for Cesium is 500 Bq/kg, which is rather conservative compared to that of Codex (1,000Bq/kg), and those of other countries.]

Discharges of radioactive substances from the Fukushima NPP into the sea have drastically decreased since early April, and to date, it is estimated that over 99% of the discharges of radioactive substances into the sea occurred during the period March 28th to April 11th. As a matter of fact, most of the recent monitoring

measurements of radioactive substances in the seawaters beyond the 30km-radius from the Fukushima NPP have constantly shown results below the detectable levels.

Nevertheless, it is reported that a number of trading companies still firmly refrain from buying fishery products from Japan, and that some governments still maintain excessive restrictions against imports of fishery products from Japan. We would like to draw your attention to and ask due consideration on the current situation and a series of monitoring efforts and control measures that the Japanese government has been taking and enforcing so as to ensure the safety of fishery products on the market.

Details

1. Monitoring programs for fishery products and restriction on fishing activities

(1) Provisional Regulatory Values in Japan

For the purpose of food safety, the Japanese government set the Provisional Regulatory Values (*) for radioactive Iodine and Cesium in fishery products at 2000 Bq/kg and 500 Bq/kg, respectively. Extensive and frequent samplings have been undertaken to ensure that no fishery products containing radioactive Iodine and/or Cesium exceeding the Provisional Regulatory Values are distributed to the market. Please note that Japan's Provisional Regulation Value for radioactive Cesium (500Bq/kg) is rather conservative compared to that of CODEX (1,000Bq/kg), as well as those of other countries (US index is 1,200Bq/kg).

[See Attachment 1: Comparison of indices for restriction on intake of foods]

(2) Monitoring programs for fishery products

The Fisheries Agency, in coordination with relevant prefectural governments, has been conducting samplings to measure levels of radioactive substances in fishery products. These samplings have been carried out at major fishing ports at least once a week for each major target species. When a measurement result detects a

level exceeding the Provisional Regulatory Values, all the related fishing activities involving that species and its landings are immediately suspended.

[See **Attachment 2**: Basic policy for Inspections on Radioactive Materials in Fishery Products]

Taking into account broad migration of some fish species, the Fisheries Agency, in close coordination with prefectural government, is undertaking samplings of fishery products in wide areas ranging from Hokkaido to Kanagawa-prefecture. The sampling measurement results are immediately posted on the websites of the Ministry of Health, Labor and Welfare (MHLW) and the Fisheries Agency. These sampling results are also made available in English on these websites.

- As of June 13, 580 samples have been measured for levels of radioactive substances, and 39 samples out of 580 showed results exceeding the Provisional Regulatory Values. Please note all these 39 samples, except for 11 samples of freshwater fish, were taken in the coastal waters close to the Fukushima NPP, consisting of limited species: epipelagic small fish (juvenile Japanese sand lance and juvenile anchovy), coastal bottom fish (Brown Hake), Invertebrates (Mediterranean mussel, Sea Urchin, Surf clam), seaweed (Wakame, Hijiki and Arame seaweed).

[See **Attachment 3**: Sampling measurement results for fishery products]

<http://www.jfa.maff.go.jp/e/inspection/index.html>

<http://www.mhlw.go.jp/english/topics/2011eq/index.html>

(3) Restriction on fishing activities and market distribution

In case where a sampling measurement detects radioactive substances exceeding the Provisional Regulatory Values, related fishing activities and landings are immediately suspended. Such suspension can be lifted only after weekly sampling measurements show constantly below the Provisional Regulatory Values three times in a row. Through such restrictive measures, no fishery products with radioactive substances exceeding the Provisional Regulatory Values are distributed to the market.

- ◆ Situation of fishing activities near Fukushima

As of June 9, the situation of fishing activities in the coastal areas in and around Fukushima is as follows:

➤ Fukushima Area

No fishing activities have been conducted since the occurrence of the Fukushima NPP accident.

➤ Miyagi area

Part of fishing activities resumed in early June, after all the sampling results of species to be caught confirmed that their levels of radioactive substances are below the Provisional Regulatory Values.

➤ Ibaraki area

Fishing activities for epipelagic fish have been suspended since sampling measurements showed that juvenile Japanese sandlances in this area exceeded the Provisional Regulatory Values. Trawl fishing resumed after all the sampling results of the species to be caught confirmed that their levels of radioactive substances are below the Provisional Regulation Values.

◆ Samplings for skipjack

Skipjack, an important species for exports from Japan, is expected to migrate soon into the offshore area of the east coast of Japan (240-320km from the coastline), and fishing activities for skipjack in that area are scheduled to begin after confirming through samplings that the levels of radioactive substances are below the Provisional Regulatory Values. During the entire fishing season, samplings are to be continuously undertaken at major fishing ports at least once a week.

2. Monitoring programs for seawaters

The Ministry of Education, Culture, Sports, Science and Technology (MEXT) and TEPCO have been conducting monitoring programs to measure the levels of radioactive substances in the seawaters and bottom sediment at over 100 sampling stations in the coastal and offshore areas in the vicinity of the Fukushima NPP. The monitoring results show a clear decreasing trend in the levels of radioactive substances. In particular, the results of the recent sampling measurements show that the levels of radioactive substances in the seawaters beyond the 30km from the Fukushima NPP have been constantly below the detectable levels*, regardless of the surface, middle, and bottom layers.

(*) The detectable levels in the monitoring by MEXT in the offshore areas are 4Bq/L

for Iodine, 6Bq/L for Cesium-134, 9Bq/L for Cesium-137. Those by TEPCO in the area around the NPP are 7Bq/L for Iodine, 15Bq/L for Cs-134 and Cs-137.

[See Attachment 4: Readings of Sea Area Monitoring (MEXT)]

<http://www.mext.go.jp/english/incident/1304192.htm>

<http://www.mext.go.jp/english/incident/1305954.htm>

Further, MEXT has conducted simulations of future diffusion and concentration of radioactive substances in seawaters, utilizing the oceanographic prediction system JCOPE-2 with oceanographic data such as ocean currents and water temperature. The recent simulation results show that radioactive substances have become and will remain below the detectable levels in the offshore areas.

[See Attachment 5: Simulation of Radioactivity Concentrations in the Sea Area (MEXT)]

<http://www.mext.go.jp/english/incident/1305758.htm>

3. For smooth transaction and exports

While the safety of fishery products on the market is secured through the monitoring efforts and restrictive measures explained above, any trade partners may require the certificates of the measurement of radioactive substances or the certificates of origin. For this, 20 inspection institutes in Japan are available to provide the measurement of radioactive substances for particular consignments of fishery products.

[See Attachment 6: List of Inspection Institutes in Japan]

In addition, the Fisheries Agency and relevant prefectural governments are currently working together to install simplified radiation measuring instruments at major fishing ports within three months so as to introduce a 100% coverage screening system for fish landings there.

[See Attachment 7: Specs of a Simplified Radiation Measuring Instrument]

4. Scientific consideration---Middle and long-term impact on fishery products

The major radioactive substances discharged from the Fukushima NPP are radioactive Iodine-131 and Cesium-134 and Cesium-137. The impact of these radioactive substances on saltwater fish is expected to be limited, considering the

following scientific facts.

(1) Dilution and diffusion of radioactive materials in the sea

Concentration levels of such radioactive substances are expected to rapidly and significantly decrease by dilution in the massive amount of the seawaters and by disperse with sea water currents/swirls. Radioactive substances released into the sea are to fall down to the bottom sediment while being attached and absorbed into suspended particles, and in the long term, they are considered to be transported to the deep-sea with the average depth of 3,800 meters, lying off the east coast of Japan.

(2) Bio-concentration of radioactive substances in saltwater fish

Radioactive half-life period of Iodine is 8 days. Therefore, even if fish intakes radioactive Iodine into its internal organs, it diminishes very shortly. For this short half-life, the transfer of radioactive iodine from seafood to human bodies is unlikely. With regard to radioactive Cesium, it has a longer half-life period (30 years). However, since Cesium behaves like Potassium in fish bodies, Cesium does not remain concentrated and is to be excreted through the gills and in the urine as the levels of radioactive substances in the surrounding seawater decreases.

The level of radioactive Cesium in fish has a close proportional relationship to that in the surrounding seawater, and it is known that the level of radioactive Cesium in fish is to decrease to around half in 50 days in the surrounding seawater with low levels of radioactive substances. This implies that measurements of radioactive substances in the seawater are important in estimating the levels of radioactive substances in fish. As mentioned above, most of the recent sampling measurements show that the levels of radioactive substances in the seawater at surface, middle, and bottom beyond the 30km-radius from the Fukushima NPP have been constantly below the limits of the detectable levels.

[See Attachment 8: Specs of a Simplified Radiation Measuring Instrument]

Attachment 1

Comparison of indices for restrictions on intake of foods

UNIT: kg/kg

	Cs-134, Cs-137						I-131		
	Drinking Water	Milk & Dairy products	Vegetables	Grain	Meat, Egg, Fish & Others	Drinking Water	Milk & Dairy products	Vegetables (ex: Root crops & Potatoes)	Others
Code	1,000	1,000	1,000	1,000	1,000	100	100	100	100
Japan	200	200	500	500	500	300	300	2,000	2,000
UA	1,200	1,200	1,200	1,200	1,200	170	170	170	170
EU	200	200	500	500	500	300	300	2,000	2,000
Thai	500	500	500	500	500	100	100	100	100
Singapore	1,000	1,000	1,000	1,000	1,000	100	100	100	100
South Korea	370	370	370	370	370	300	150	300	300
Hong Kong	1,000	1,000	1,000	1,000	1,000	100	100	100	100
Chinese Taipei	370	370	370	370	370	300	55	300	300
Philippines	1,000	1,000	1,000	1,000	1,000	1,000	1,000	1,000	1,000
Vietnam	1,000	1,000	1,000	1,000	1,000	100	100	100	100
Malaysia	1,000	1,000	1,000	1,000	1,000	100	100	100	100
China		300	210	200	Meat, Fish & Crustacean 100, Poultry 100, Potatoes 30			160	Meat and Fishery Product 100, Eggs 150, Potatoes 60

Attachment 2

(Provisional translation)

Press Release

6 May, 2011

Fisheries Agency of Japan

The Fisheries Agency of Japan has established the "Basic Policy for Inspections on Radioactive Materials in Fishery Products", and notified to the relevant Prefectural and Metropolitan Governments and organizations.

Basic Policy for Inspections on Radioactive Materials in Fishery Products

Inspections on radioactive materials in fishery products have been conducted in response to the emission of radioactive materials from the TEPCO Fukushima Daiichi Nuclear Power Plant. In order to strengthen our inspection on radioactive materials in fishery products, including fish species with their peak fishing season in upcoming months, the Fisheries Agency of Japan has established the "Basic Policy for Inspections on Radioactive Materials in Fishery Products", as follows.

1. Basic policy for inspection

(1) Inspection of coastal species

(a) From Kanagawa Prefecture to the southern part of Fukushima Prefecture

Based on formation of fishing grounds off the coast of each prefecture, Prefectural Governments should designate areas where inspection is necessary, and conduct sampling once a week in principle (once every two weeks in Kanagawa Prefecture and islands belonging to Tokyo Metropolis), at the main landing ports of each designated area. When sampling is conducted at markets, the area where the fish was caught should be confirmed.

Major species caught in each fishing season should be selected as the target species for inspection, taking into local circumstances into account. They should be selected to cover a wide spectrum of marine habitat such as surface (e.g. Juvenile Japanese sand lance), middle column (e.g. sea bass, sea bream) and bottom (e.g. flounder, conger eel), taking into account that larger amount of radioactive materials has been detected so far in species swimming in the surface (sand lance).

(b) Northern part of Fukushima Prefecture and to the north

Inspection should be conducted before resumption of fishery operations. Decision on whether to resume fishery operations should be based on the analysis results of the inspection.

When fishery operation is resumed, Prefectural Governments should designate areas where inspection is necessary, and conduct sampling once a week in principle (once every two weeks in Iwate Prefecture and to the north), at the main landing ports of each designated area.

Target species for inspection should be selected in accordance with the procedure (a) above.

(2) Migratory species (skipjack, Japanese jack mackerel, Pacific saury, etc.)

Inspections should be conducted through cooperation between relevant fisheries industry organizations and the Prefectural Governments where the fish is landed. (The inspection framework when the fishing grounds of such fish species move northward is under discussion among relevant parties, including industry organizations. Use of Hokusho-maru, a research fishing vessel of the Federation of North Pacific District Purse Seine Fisheries Co-operative Associations of Japan, in this framework will be considered.)

(a) Skipjack

After the formulation of fishing grounds off the coast of Izu Islands and Boso Peninsula (around the middle of May), inspections should be conducted once a week in principle (sampling should be conducted at the fishing ports in Chiba Prefecture where landing of skipjack is expected (namely Choshi and Katsuura fishing ports)).

When formation of fishing grounds off the coast of Fukushima Prefecture (usually 240-320 km off the coast) is expected (around early June), sampling by a trial fishing vessel should be conducted prior to commercial operations. Decision on whether to operate fishery in the area should be based on the analysis results. When fishery operation is to continue, sampling should be conducted once a week in principle at landing ports.

When fishing grounds are formed off the coast of Miyagi Prefecture and to the north, inspections should be conducted once a week in principle.

(b) Sardine and mackerel

While fishing grounds are formed off the coast of Chiba Prefecture, sampling should continue at the fishing ports in Chiba Prefecture where landing of sardine and mackerel is expected (namely Choshi fishing port).

When formation of fishing grounds off the coast of Ibaraki Prefecture is expected (in May), sampling by the research vessel of the Ibaraki Prefectural Fisheries Experimental Station should be conducted, in cooperation with the Ibaraki Prefectural Government. Decision on whether to operate fishery should be based on the analysis results. When fishery operation is to continue, sampling should be conducted once a week in principle at landing ports.

When formation of fishing grounds off the coast of Fukushima Prefecture is expected (in June), sampling should be conducted by a research fishing vessel. The rest of the procedure will be the same as described above.

When fishing grounds are formed off the coast of Miyagi Prefecture and to the north, inspections should be conducted once a week in principle.

(c) Pacific saury and salmon migrating southward

Starting from summer, inspections should be conducted once a week in principle.

2. Amount of sample

Sample size should be a sufficient amount to conduct inspection; i.e. 5kg or more per species in principle. The sampling site and date should be recorded.

3. Additional notes

Due to the migratory nature of fish, and to varying weather conditions, sampling of target species at the scheduled site and date may not always be possible. Sampling plans should be drawn up with ample flexibility to allow for these conditions.

4. Publication of inspection results

The publication and reporting to the Ministry of Health, Labor and Welfare of inspection results should be conducted by the prefectural Government in whose water the sample is caught, or in which the sampling port is located.

5. Response to inspection results that exceed the Provisional Regulation Value in migratory species

When inspection results exceeding the Provisional Regulation Value are detected in migratory species, the industry concerned will be requested to voluntarily refrain from relevant fishing operations around the site where the sample was caught (generally on a prefecture by prefecture basis). Then sampling by a research fishing vessel should be

conducted once a week in principle. Fishery operations could resume only after the inspection results are below the Provisional Regulation Value for 3 consecutive times.

For more information
Resources and Environment Research Division
Fisheries Agency of Japan
+81-3-3502-8487

Attachment 3

Fisheries Agency of Japan

Summary of Monitoring Results of the levels of Radioactive Substances in Fishery products (As of June 13, 2011)

As of June 13, 580 fishery products have been sampled for the measurement the levels of radioactive materials. 39 samples out of the 580 showed the results exceeding the Provisional Regulatory Values.

Saltwater fish	333 (*18)	<div>Juvenile Japanese sandlance(12) White bait(4) Fat greenling (1) Brown Hake(1)</div>
Invertebrates	85 (*5)	<div>Mediterranean mussel(1) Sea urchin(2) Surf clam(2)</div>
Seaweed	26 (*5)	<div>Wakame seaweed(1) Hijiki seaweed(1) Arame seaweed(3)</div>
Seafood products	14 (*0)	
Freshwater fish	117 (*11)	<div>Ayu sweetfish(3) Land-locked salmon(4) Japanese smelt(2) Japanese dace (2)</div>
Marine Mammals	5 (*0)	

(*) Figures in brackets show the number of samples with results exceeding the Provisional Regulatory Values

Implementation of the inspection on radioactivity level in fisheries products (No.1)


As of 7 June

All fisheries activities are voluntarily refrained in Fukushima Prefecture.


Fish species that exceeded the Provisional Regulation Value are Japanese sand lance, whitebait, ayu sweetfish, Japanese smelt, Land-locked salmon, Mediterranean mussel, Wakame seaweed, Hijiki seaweed, Arame seaweed, Japanese dace, Surf clam and Sea urchin

2011 Fishing season for Japanese sand lance has ended in all the Ibaraki Prefecture area.

[Note 1]
● Indicate the inspection results exceeding the Provisional Regulation Value.
○ Indicate the inspection results below the Provisional Regulation Value




Implementation of the inspection on radioactivity level
in fisheries products (No.2)




Attachment 4

海域モニタリング結果(平成23年5月23日~27日採水)
Readings of Sea Area Monitoring (May 23-27, 2011)



海域モニタリング結果(平成23年5月21日～22日採水)
Readings of Sea Area Monitoring (May 21-22, 2011)



上記測定点の海水温及び塩分濃度については、独立行政法人 海洋研究開発機構の下記Webにて公開している。
The readings of temperatures and salinity levels of seawater at the measurement points are put on the websites of JAMSTEC below.
<http://www.godac.jamstec.go.jp/monitoringdata/>


東京電力株式会社福島第一原子力発電所周辺の海水中の放射能濃度分布

(Distribution map of radioactivity concentration in the seawater

around TEPCO Fukushima Dai-ichi NPP)

(試料採取日:平成23年6月11日)

(Sampling Date: 2011/6/11)



図中の■は東京電力福島第一発電所を示す


※東京電力(株)の発表(<http://www.tepco.co.jp/oc/press/index11-j.html>)をもとに文部科学省が作成

(Based on the press release of TEPCO (<http://www.tepco.co.jp/oc/press/index11-j.html>))

※1 NDの記載は、海水の放射能濃度の検出値が検出限界値(I-131が約6Bq/L、Cs-134が約14Bq/L及びCs-137が約15Bq/L)を下回る場合。

※1 ND indicates the case that the detected radioactivity concentration in sea water was lower than the detection limits of approximately 6 Bq/L for I-131, 14 Bq/L for Cs-134 and 15 Bq/L for Cs-137.

東京電力株式会社福島第一原子力発電所周辺の海水中の放射能濃度分布
 (Distribution map of radioactivity concentration in the seawater
 around TEPCO Fukushima Dai-ichi NPP)
 (試料採取日:平成23年6月10日)
 (Sampling Date: 2011/6/10)



図中の■は東京電力福島第一発電所を示す

※東京電力(株)の発表(<http://www.tepco.co.jp/cs/press/index11-j.html>)をもとに文部科学省が作成

(Based on the press release of TEPCO (<http://www.tepco.co.jp/cs/press/index11-j.html>))

※1 NDの記載は、海水の放射能濃度の検出値が検出限界値(I-131が約7Bq/L、Cs-134が約14Bq/L及びCs-137が約15Bq/L)を下回る場合。

※2 ND indicates the case that the detected radioactivity concentration in sea water was lower than the detection limits of approximately 7 Bq/L for I-131, 14 Bq/L for Cs-134 and 15 Bq/L for Cs-137.

東京電力株式会社福島第一原子力発電所周辺及び茨城県沿岸の海水中の放射能濃度分布

(Distribution map of radioactivity concentration in the seawater


around TEPCO Fukushima Dai-ichi NPP and coast of Ibaraki Prefecture)

(試料採取日: 茨城県沿岸 平成23年6月7日～8日)

(Sampling Date: Coast of Ibaraki Prefecture 2011/6/7~2011/6/8)

(試料採取日: 東京電力福島第一原子力発電所周辺 平成23年6月9日)

(Sampling Date: Around TEPCO Fukushima Dai-ichi NPP 2011/6/9)



図中の■は東京電力福島第一発電所を示す

*東京電力(株)の発表(<http://www.tepco.co.jp/oc/press/index11-j.html>)をもとに文部科学省が作成

(Based on the press release of TEPCO (<http://www.tepco.co.jp/oc/press/index11-j.html>))


※1 NDの記載は、海水の放射能濃度の検出値が検出限界値(東京電力福島第一発電所周辺についてはI-131が約68Bq/L、Cs-134が約148Bq/L及びCs-137が約158Bq/L、茨城県沿岸についてはI-131が約8Bq/L、Cs-134が約15Bq/L及びCs-137が約138Bq/L)を下回る場合。

※1 ND indicates the case that the detected radioactivity concentration in sea water was lower than the detection limits of approximately 6 Bq/L for I-131, 14 Bq/L for Cs-134 and 15 Bq/L for Cs-137 for around TEPCO Fukushima Dai-ichi NPP, approximately 8 Bq/L for I-131, 15 Bq/L for Cs-134 and 12 Bq/L for Cs-137 for coast of Ibaraki Prefecture.


Attachment 5

[[Figure 3-1] Simulation of Radioactivity Concentrations by JCOPE2 (May 31)
(based on data up to May 20)

JCOPE2 Predicted figures (I 137)
2011/05/31



JCOPE2 Predicted figures (I 134)
2011/05/31




Below the detection limit values

(Note: The index numbers in the figure show how many times the maximum radioactivity concentrations predicted at each point are higher than the effluent concentration limits for nuclear facilities determined by regulations (Cs134: 60Bq/L, Cs137: 90Bq/L).)

*Currently, detection limit values are 6Bq/L(Cs134) and 9Bq/L(Cs137) in monitoring of sea areas near Fukushima Dai-ichi NPP by the Ministry of Education, Culture, Sports, Science and Technology (MEXT).

[[Figure 3-2] Simulation of Radioactivity Concentrations by JCOPE2 (June 15)
(based on data up to May 20)




Below the detection limit values

{Note: The index numbers in the figure show how many times the maximum radioactivity concentrations predicted at each point are higher than the effluent concentration limits for nuclear facilities determined by regulations (Cs134: 60Bq/L, Cs137: 90Bq/L).}

*Currently, detection limit values are 6Bq/L(Cs134) and 9Bq/L(Cs137) in monitoring of sea areas near Fukushima Dai-ichi NPP by the Ministry of Education, Culture, Sports, Science and Technology (MEXT).

[Figure 3-3] Simulation of Radioactivity Concentrations by JCOPE2 (June 30)
(based on data up to May 20)




Below the detection limit values

(Note: The index numbers in the figure show how many times the maximum radioactivity concentrations predicted at each point are higher than the effluent concentration limits for nuclear facilities determined by regulations (Cs134: 60Bq/L, Cs137: 90Bq/L).)

*Currently, detection limit values are 6Bq/L (Cs134) and 9Bq/L (Cs137) in monitoring of sea areas near Fukushima Dai-ichi NPP by the Ministry of Education, Culture, Sports, Science and Technology (MEXT).

[Figure 3-4] Simulation of Radioactivity Concentrations by JCOPE2 (July 15)
(based on data up to May 20)



(Note: The index numbers in the figure show how many times the maximum radioactivity concentrations predicted at each point are higher than the effluent concentration limits for nuclear facilities determined by regulations (Cs134: 60Bq/L, Cs137: 90Bq/L).)

*Currently, detection limit values are 6Bq/L (Cs134) and 9Bq/L (Cs137) in monitoring of sea areas near Fukushima Dai-ichi NPP by the Ministry of Education, Culture, Sports, Science and Technology (MEXT).


List of Inspection Institutes for Radioactive Substances in Foods

No.	Inspection Institute (Japanese)	Inspection Institute (English)	Inspection Equipment	Status
1	京都市衛生環境研究所	-	Germanium Semiconductor Detector	In operation
2	(財)食品環境検査協会	Japan Inspection Association of Food and Food Industry Environment	Germanium Semiconductor Detector	In operation
3	(財)日本分析センター	Japan Chemical Analysis Center	Germanium Semiconductor Detector	In operation
4	(財)日本冷凍食品検査協会	Japan Frozen Foods Inspection Corporation	Germanium Semiconductor Detector	In operation
5	(財)日本食品分析センター	Japan Food Research Laboratories	Germanium Semiconductor Detector	In operation
6	(社)青森県薬剤師会衛生検査センター	-	NaI Scintillation Counter	In operation
7	(財)千葉県薬剤師会検査センター	-	Germanium Semiconductor Detector	In operation
8	(財)日本乳業技術協会	Japan Dairy Technical Association	Germanium Semiconductor Detector	In operation
9	(財)新日本検定協会	Shin Nihon Kentei Kyokai	Germanium Semiconductor Detector	In operation
10	(財)北海道薬剤師会公衆衛生検査センター	-	NaI Scintillation Counter	In operation
11	(財)新潟県環境衛生研究所	Environmental Science Research Niigata	Germanium Semiconductor Detector	In operation
12	(社)新潟県環境衛生中央研究所	Niigata Environment Hygiene Central Laboratory Co.	Germanium Semiconductor Detector	In operation
13	(社)日本食品衛生協会食品衛生研究所	Japan Food Hygiene Association Institute of Food Hygiene	Germanium Semiconductor Detector	In operation
14	(一般社団法人)日本海事検定協会	Nippon Kaiji Kentei Kyokai	Germanium Semiconductor Detector	Start in mid-June
15	(株)島津テクノリサーチ	Shimadzu Techno-Research	Germanium Semiconductor Detector	Start in early-July
16	(財)食品分析開発センター-SUNATEC	Food Analysis Technology Center	Germanium Semiconductor Detector	Start in early-July
17	(株)日本食品エコロジ-研究所	Japan Institute of Foods Ecology	Germanium Semiconductor Detector	Start in early-July
18	(株)北陸環境科学研究所	-	Germanium Semiconductor Detector	Start in early-July
19	(株)神環検査センター	Seikan	Germanium Semiconductor Detector	Start in early-July
20	(財)東京環境検院	-	Germanium Semiconductor Detector	Start in late-July

Attachment 6

Attachment 7

ACTIVITY MEASUREMENT



Activity Monitor LB 2045/LB 742

Modern gamma spectroscopy system for nuclide-specific activity measurements: activity in foodstuffs, liquids, bulk goods in Bq/l

- Detection limit approx. 1 Bq/l
- Connection for scintillator probes
- 3 different energy ranges
- Spectrum presentation (1024 channels)
- One or two energy windows
- Nuclide library
- Storage of spectra or ROI data
- Half-life correction
- Weight entry and spillover correction.

Intuitive user guidance: operation via softkeys on the display

Several service functions are available: background measurement, energy calibration and spectrum recording

Measurement/Spectrum

ADC	1024 channels, 7 µs max. conversion time
Energy ranges	0-256 keV, 0-1024 keV, 0-2048 keV
Energy calibration	non-linear amplifical function
Region of Interest	max. 2 ROIs
Nuclide library	max. 50 nuclides

Data Acquisition/Computer

Processor	Motorola MC88340 32 bit 16 MHz
Display/Touch panel	Graphical LC display 320 x 240 pixels
Memory	max. 70 spectra or 800 ROI values
Serial interface	RS232

General Specifications

High voltage supply	0-1300 Volt, polarity positive, resolution 12 bit
Power supply (wide-range input)	85-264 VAC, 47-65 Hz
Protection class	IP 54
Temperature range	-5°C to +40°C
Relative humidity	0% to 90% (no condensation)
External dimensions	245 x 145 x 325 mm (W x H x T)
Weight	3.3 kg

SZ 50 U 6S8/2E-X scintillation probe

Detector	Nal crystal 2" x 2", with photomultiplier and voltage divider
Resolution	7.5 % (FWHM) for ¹³⁷ Cs 661 keV
External diameter	55 mm
Complete cable set	2 m

Shielding LB 7428 A

Lead chamber for 2" probes	
50 mm lead shielding for 0.4 l Marinelli beakers	
Weight	approx. 75 kg

日本代理店

ベルトールドジャパン株式会社


東京本社
〒111-0052 東京都台東区根岸1-9-1 根岸ティールビル3F
Tel.03-5825-3557 Fax.03-5825-3558
大阪営業所
〒532-0004 大阪府大阪市淀川区西宮原1-4-25 第2番ビル4F
Tel.06-6393-5551 Fax.06-6393-3331
<http://www.berthold-jp.com>

Bio-accumulation or bio-concentration of radionuclides through food chain

$$\text{Concentration factor} = \frac{\text{Concentration in fish body}}{\text{Concentration in sea water}}$$

Concentration Factor of marine fish

Materials	Concentration Factor
Cs	5 ~ 100
I	10
U	10
Pt	3.5
Hg	360 ~ 600
DDT	1200
PCB	1200 ~ 10000



* Bio-accumulation or bio-concentration of radionuclides through food chain is not increasing.

Why are not accumulated ?

* Very low Concentration Factors

Reference:
Fujio Kasamatsu
bio-concentration Edit. N. Yamagata,
Radioisotopes 48, 1999.




Iodine and Cesium

- Iodinesolid/gaseous (sublimation nucleotide)
I-131 (Half life time: 8.04 days)
- Cs.....solid , behaves like potassium :
does not accumulate to specific organs
Cs-137 (Half life time :30.1years) ,
Cs-134 (Half life time: 2.07years)

Periodic table

	1A	2A	3A	4A	5A	6A	7A	8	1B	2B	3B	4B	5B	6B	7B	0
1	H															He
2	Li	Be														Ne
3	Na	Mg														Ar
4	K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn				Kr
5	Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd				Xe
6	Cs	Ba	*	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg				Rn
7	Fr	Ra	**													
*	La	Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu	
**	Ac	Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr	


The flow of salts in marine fish body



• Radionuclides excrete, not accumulate.

• The concentration in fish is depend on the concentration of environmental water .

Comparison of Cs-137 concentration between sea water and fish body





Cs-137 concentration annual changes in Japan coast

- Fish body concentration depends on sea water concentration



Excretion of radio nucleotides



• In natural condition
 Cs-137 excretes
quickly.

Ref.
K. Yoshida, JCAC 34, 1999.
F. Kasamatsu, Radiotopes 48,
1999.

東京電力株式会社福島第一原子力発電所周辺及び茨城県沿岸の海水中の放射能濃度分布

(Distribution map of radioactivity concentration in the seawater


around TEPCO Fukushima Dai-ichi NPP and coast of Ibaraki Prefecture)

(試料採取日: 茨城県沿岸 平成23年6月7日~8日)

(Sampling Date: Coast of Ibaraki Prefecture 2011/6/7~2011/6/8)

(試料採取日: 東京電力福島第一原子力発電所周辺 平成23年6月9日)

(Sampling Date: Around TEPCO Fukushima Dai-ichi NPP 2011/6/9)



図中の■は東京電力福島第一発電所を示す

*東京電力(株)の発表(<http://www.tepco.co.jp/ci/press/index11-j.html>)をもとに文部科学省が作成

(Based on the press release of TEPCO (<http://www.tepco.co.jp/ci/press/index11-j.html>))


※1 NDの記載は、海水の放射能濃度の検出値が検出限界値(東京電力福島第一発電所周辺についてはI-131が約60Bq/L、Cs-134が約14Bq/L及びCs-137が約15Bq/L、茨城県沿岸についてはI-131が約8Bq/L、Cs-134が約16Bq/L及びCs-137が約13Bq/L)を下回る場合。

※1 ND indicates the case that the detected radioactivity concentration in sea water was lower than the detection limits of approximately 6 Bq/L for I-131, 14 Bq/L for Cs-134 and 15 Bq/L for Cs-137 for around TEPCO Fukushima Dai-ichi NPP, approximately 8 Bq/L for I-131, 16 Bq/L for Cs-134 and 13 Bq/L for Cs-137 for coast of Ibaraki Prefecture.

Attachment 5

[Figure 3-1] Simulation of Radioactivity Concentrations by JCOPE2 (May 31)
(based on data up to May 20)


JCOPE2 Predicted figures (I 134)
2011/05/31



Below the detection limit values


(Note: The index numbers in the figure show how many times the maximum radioactivity concentrations predicted at each point are higher than the effluent concentration limits for nuclear facilities determined by regulations (Cs134: 60Bq/L, Cs137: 90Bq/L).)
*Currently, detection limit values are 6Bq/L (Cs134) and 9Bq/L (Cs137) in monitoring of sea areas near Fukushima Dai-ichi NPP by the Ministry of Education, Culture, Sports, Science and Technology (MEXT).

JCOPE2 Predicted figures (I 137)
2011/05/31



Below the detection limit values

[[Figure 3-2] Simulation of Radioactivity Concentrations by JCOPE2 (June 15)
(based on data up to May 20)]




Below the detection limit values

(Note: The index numbers in the figure show how many times the maximum radioactivity concentrations predicted at each point are higher than the effluent concentration limits for nuclear facilities determined by regulations (Cs134: 60Bq/L, Cs137: 90Bq/L).)

*Currently, detection limit values are 6Bq/L(Cs134) and 9Bq/L(Cs137) in monitoring of sea areas near Fukushima Dai-ichi NPP by the Ministry of Education, Culture, Sports, Science and Technology (MEXT).

[(Figure 3-3] Simulation of Radioactivity Concentrations by JCOPE2 (June 30)
(based on data up to May 20)




Below the detection limit values

Below the detection limit values

(Note: The index numbers in the figure show how many times the maximum radioactivity concentrations predicted at each point are higher than the effluent concentration limits for nuclear facilities determined by regulations (Cs134: 60Bq/L, Cs137: 90Bq/L).)

*Currently, detection limit values are 6Bq/L(Cs134) and 9Bq/L(Cs137) in monitoring of sea areas near Fukushima Dai-ichi NPP by the Ministry of Education, Culture, Sports, Science and Technology (MEXT).

[[Figure 3-4] Simulation of Radioactivity Concentrations by JCOPE2 (July 15)
(based on data up to May 20)



{Note: The index numbers in the figure show how many times the maximum radioactivity concentrations predicted at each point are higher than the effluent concentration limits for nuclear facilities determined by regulations (Cs134: 60Bq/L, Cs137: 90Bq/L).}

*Currently, detection limit values are 6Bq/L(Cs134) and 9Bq/L(Cs137) in monitoring of sea areas near Fukushima Dai-ichi NPP by the Ministry of Education, Culture, Sports, Science and Technology (MEXT).


List of Inspection Institutes for Radioactive Substances in Foods

No.	Inspection Institute (Japanese)	Inspection Institute (English)	Inspection Equipment	Status
1	京都市衛生環境研究所	-	Germanium Semiconductor Detector	In operation
2	(財)食品環境検査協会	Japan Inspection Association of Food and Food Industry Environment	Germanium Semiconductor Detector	In operation
3	(財)日本分析センター	Japan Chemical Analysis Center	Germanium Semiconductor Detector	In operation
4	(財)日本冷凍食品検査協会	Japan Frozen Foods Inspection Corporation	Germanium Semiconductor Detector	In operation
5	(財)日本食品分析センター	Japan Food Research Laboratories	Germanium Semiconductor Detector	In operation
6	(社)青森県薬剤師会衛生検査センター	-	NaI Scintillation Counter	In operation
7	(財)千葉県薬剤師会検査センター	-	Germanium Semiconductor Detector	In operation
8	(財)日本乳業技術協会	Japan Dairy Technical Association	Germanium Semiconductor Detector	In operation
9	(財)新日本検定協会	Shin Nihon Kentei Kyokai	Germanium Semiconductor Detector	In operation
10	(財)北海道薬剤師会公衆衛生検査センター	-	NaI Scintillation Counter	In operation
11	(財)新潟県環境衛生研究所	Environmental Science Research Niigata	Germanium Semiconductor Detector	In operation
12	(社)新潟県環境衛生中央研究所	Niigata Environment Hygiene Central Laboratory Co.	Germanium Semiconductor Detector	In operation
13	(社)日本食品衛生協会食品衛生研究所	Japan Food Hygiene Association Institute of Food Hygiene	Germanium Semiconductor Detector	In operation
14	(一般社団法人)日本海軍検定協会	Nippon Kaiji Kentei Kyokai	Germanium Semiconductor Detector	Start in mid-June
15	(株)島津テクノリサーチ	Shimadzu Techno-Research	Germanium Semiconductor Detector	Start in early-July
16	(財)食品分析開発センター-SUNATEC	Food Analysis Technology Center	Germanium Semiconductor Detector	Start in early-July
17	(株)日本食品エコロジ-研究所	Japan Institute of Foods Ecology	Germanium Semiconductor Detector	Start in early-July
18	(株)北陸環境科学研究所	-	Germanium Semiconductor Detector	Start in early-July
19	(株)静環検査センター	Seikan	Germanium Semiconductor Detector	Start in early-July
20	(財)東京環境検院	-	Germanium Semiconductor Detector	Start in late-July

Attachment 6

Attachment 7

ACTIVITY MEASUREMENT



Activity Monitor LB 2045/LB 742

Modern gamma spectroscopy system for nuclide-specific activity measurements activity in foodstuffs, liquids, bulk goods in Bq/l

- Detection limit approx. 1 Bq/l
- Connection for scintillator probes
- 3 different energy ranges
- Spectrum presentation (1024 channels)
- One or two energy windows
- Nuclide library
- Storage of spectra or ROI data
- Half-life correction
- Weight entry and spillover correction.

Intuitive user guidance: operation via softkeys on the display

Several service functions are available: background measurement, energy calibration and spectrum recording

Measurement/Spectrum

ADC	1024 channels, 7 μ s max. conversion time
Energy ranges	0-256 keV, 0-1024 keV, 0-2048 keV
Energy calibration	non-linear empirical function
Region of Interest	max. 2 ROIs
Nuclide library	max. 50 nuclides

Data Acquisition/Computer

Processor	Motorola MC68340 32 bit 16 MHz
Display/Touch panel	Graphical LC display 320 x 240 pixels
Memory	max. 70 spectra or 800 ROI values
Serial interface	RS232

General Specifications

High voltage supply	0-1300 Volt, polarity positive, resolution 12 bit
Power supply (wide-range input)	85-264 VAC, 47-65 Hz
Protection class	IP 54
Temperature range	-5°C to +40°C
Relative humidity	0% to 90% (no condensation)
External dimensions	245 x 145 x 325 mm (W x H x T)
Weight	3.3 kg

SZ 50 U 8S8/2E-X scintillation probe

Detector	NaI crystal 2" x 2", with photomultiplier and voltage divider
Resolution	7.5 % (FWHM) for ^{137}Cs 661 keV
External diameter	55 mm
Complete cable set	2 m

Shielding LB 7428 A

Lead chamber for 2" probes	
50 mm lead shielding for 0.4 l Marinelli beakers	
Weight	approx. 75 kg


日本代理店

ベルトールドジャパン株式会社

東京本社
〒111-0052 東京都台東区根岸1-9-1 郵便ティールビル3F
Tel.03-5625-3557 Fax.03-5625-3558

大阪営業所
〒532-0004 大阪府大阪市淀川区西宮西1-4-25 第2ビル4F
Tel.06-6393-5551 Fax.06-6393-3331
<http://www.berthold-jp.com>

Bio-accumulation or bio-concentration of radionuclides through food chain



• Bio-accumulation or bio-concentration of radionuclides through food chain is not increasing.

Why are not accumulated ?¹

• Very low Concentration Factors

Reference:
 Fujio Kasamatsu
 bio-concentration Edit. N. Yamagata,
 Radioisotopes 48, 1999.




Iodine and Cesium

- Periodic table


	1A	2A	3A	4A	5A	6A	7A	8	1B	2B	3B	4B	5B	6B	7B	0
1	H															He
2	Li	Be														Ne
3	Na	Mg														Ar
4	K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn				Kr
5	Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd				Xe
6	Cs	Ba	*	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg				Rn
7	Fr	Ra	**													
*	La	Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu	
**	Ac	Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr	

The flow of salts in marine fish body



- Radionucleotides excrete, not accumulate.
- The concentration in fish is depend on the concentration of environmental water.

Comparison of Cs-137 concentration between sea water and fish body





Cs-137 concentration annual changes in Japan coast

- Fish body concentration depends on sea water concentration



Excretion of radio nucleotides



• In natural condition Cs-137 excretes quickly.

Ref. K. Yoshida, JCAC 34, 1999.
F. Kasamatsu, Radiotopes 48, 1999.